

# Prethermalization of quantum systems interacting with non-equilibrium environments

New J. Phys. 22 083067

Andreu Anglés-Castillo<sup>1</sup>, Mari Carmen Bañuls<sup>2,3</sup>,  
Armando Pérez<sup>1</sup>, Inés de Vega<sup>4</sup>

<sup>1</sup>Department of Theoretical Physics, UV - IFIC, Valencia, Spain

<sup>2</sup>Max-Planck-Institut für Quantenoptik, Hans-Kopfermann-Str. 1, 85748 Garching, Germany

<sup>3</sup>MCQST, Schellingstr. 4, D-80799 München, Germany

<sup>4</sup>Department of Physics and Arnold Sommerfeld Center for Theoretical Physics,  
Ludwig-Maximilians-University Munich, Germany

May 11, 2021

# Model




Total Hamiltonian:

$$H = H_S$$

Components:

$$H_S = \frac{1}{2}\omega_0\sigma_z$$

# Model



Total Hamiltonian:


$$H = H_S + H_I + H_{RI}$$

Components:

$$H_S = \frac{1}{2}\omega_0\sigma_z$$

$$H_{RI} = \sum_{\lambda} \omega_{\lambda} a_{\lambda}^{\dagger} a_{\lambda} \quad H_I = \sum_{\lambda} g_{\lambda} \sigma_{+} a_{\lambda} + h.c. ,$$

# Model



Total Hamiltonian:

$$H = H_S + H_I + H_{RI} + H_{I2} + H_{RII}$$

Components:

$$H_S = \frac{1}{2}\omega_0\sigma_z$$

$$H_{RI} = \sum_{\lambda} \omega_{\lambda} a_{\lambda}^{\dagger} a_{\lambda} \quad H_I = \sum_{\lambda} g_{\lambda} \sigma_{+} a_{\lambda} + h.c. ,$$

$$H_{RII} = \sum_{\lambda,k} \omega_{\lambda,k} b_{\lambda,k}^{\dagger} b_{\lambda,k} \quad H_{I2} = \sum_{\lambda,k} g_{\lambda,k} a_{\lambda} b_{\lambda,k}^{\dagger} + h.c.$$

# Model

Some further assumptions:

- ▶ All the reservoirs around each single mode of the first reservoir are identical

# Model

Some further assumptions:

- ▶ All the reservoirs around each single mode of the first reservoir are identical
- ▶ The initial state of system + environment is a product state

$$\rho(0) = \rho_S(0) \otimes \rho_E$$

# Model

Some further assumptions:

- ▶ All the reservoirs around each single mode of the first reservoir are identical
- ▶ The initial state of system + environment is a product state

$$\rho(0) = \rho_S(0) \otimes \rho_E$$

- ▶ The different reservoirs are also in a product state of thermal states

$$\rho_E = \rho_{th}(\beta_I) \otimes \rho_{th}(\beta_{II}) \quad \rho_{th}(\beta_i) \propto e^{-H_{Ri}\beta_i}$$

# Model

Some further assumptions:

- ▶ All the reservoirs around each single mode of the first reservoir are identical
- ▶ The initial state of system + environment is a product state

$$\rho(0) = \rho_S(0) \otimes \rho_E$$

- ▶ The different reservoirs are also in a product state of thermal states

$$\rho_E = \rho_{th}(\beta_I) \otimes \rho_{th}(\beta_{II}) \quad \rho_{th}(\beta_i) \propto e^{-H_{Ri}\beta_i}$$

- ▶ We model the spectral functions of the reservoirs with the phenomenological model:

$$J_i(\omega) = g_i \omega^{s_i} e^{-\omega/\omega_c}, \quad i = \text{RI, RII}$$

# Methods

## Redfield Master Equation

$$\frac{d}{dt}\rho_S(t) = -i[H_S, \rho_S(t)] + \left( \int_0^t d\tau \alpha^+(t, \tau) [V_{\tau-t} \sigma_+ \rho_S(t), \sigma_-] + \int_0^t d\tau \alpha^-(t, \tau) [V_{\tau-t} \sigma_- \rho_S(t), \sigma_+] + h.c. \right),$$

where  $V_t \mathcal{O} = e^{iH_S t} \mathcal{O} e^{-iH_S t}$  represents the free evolution of the operator.

# Methods

## Canonical form of the Master Equation

$$\frac{d}{dt}\rho_S(t) = -i[H(t), \rho_S(t)] + \sum_{k=1}^{d^2-1} \gamma_k(t) \left( L_k(t)\rho_S(t)L_k^\dagger(t) - \frac{1}{2}\{L_k^\dagger(t)L_k(t), \rho_S(t)\} \right),$$

# Methods

## Canonical form of the Master Equation

$$\frac{d}{dt}\rho_S(t) = -i[H(t), \rho_S(t)] + \sum_{k=1}^{d^2-1} \gamma_k(t) \left( L_k(t)\rho_S(t)L_k^\dagger(t) - \frac{1}{2}\{L_k^\dagger(t)L_k(t), \rho_S(t)\} \right),$$

with

$$\gamma_+(t) = J_I(\omega_0)n_I(\omega_0)e^{-J_{II}(\omega_0)t} + J_I(\omega_0)n_{II}(\omega_0)(1 - e^{-J_{II}(\omega_0)t}), \quad L_+ = \sigma_+$$

$$\gamma_-(t) = J_I(\omega_0)(n_I(\omega_0)+1)e^{-J_{II}(\omega_0)t} + J_I(\omega_0)(n_{II}(\omega_0)+1)(1 - e^{-J_{II}(\omega_0)t}), \quad L_- = \sigma_-$$

where

$$n_i(\omega) = \frac{1}{e^{-\beta_i\omega} - 1}$$

## Results - Prethermalization

This model exhibits an instance of prethermalization: **relaxation of the open system to a state that is not the true asymptotic state**

# Results - Prethermalization

This model exhibits an instance of prethermalization: **relaxation of the open system to a state that is not the true asymptotic state**

When present, we can differentiate the following steps:

1. Relaxation of any initial condition to a thermal state determined by the temperature of RI.

## Results - Prethermalization

This model exhibits an instance of prethermalization: **relaxation of the open system to a state that is not the true asymptotic state**

When present, we can differentiate the following steps:

1. Relaxation of any initial condition to a thermal state determined by the temperature of RI.
2. The system remains stationary in that state

## Results - Prethermalization

This model exhibits an instance of prethermalization: **relaxation of the open system to a state that is not the true asymptotic state**

When present, we can differentiate the following steps:

1. Relaxation of any initial condition to a thermal state determined by the temperature of RI.
2. The system remains stationary in that state
3. Final relaxation towards a thermal state determined by the temperature of RII.

# Results - Prethermalization

Animations!!

## Result - Prethermalization time

Trace distance between the prethermal state and the evolved state of the system to study dependence of prethermalization time on other parameters.

$$T(\rho_S(t), \rho_S^{\text{th}}(\beta_I)) = \frac{1}{2} \text{Tr} \left\{ \sqrt{(\rho_S(t) - \rho_S^{\text{th}}(\beta_I))^2} \right\} \quad (1)$$

## Result - Prethermalization time

Trace distance between the prethermal state and the evolved state of the system to study dependence of prethermalization time on other parameters.

$$T(\rho_S(t), \rho_S^{\text{th}}(\beta_I)) = \frac{1}{2} \text{Tr} \left\{ \sqrt{(\rho_S(t) - \rho_S^{\text{th}}(\beta_I))^2} \right\} \quad (1)$$

- ▶ Exponential dependence with the coupling strength between environments.

## Result - Prethermalization time

Trace distance between the prethermal state and the evolved state of the system to study dependence of prethermalization time on other parameters.

$$T(\rho_S(t), \rho_S^{\text{th}}(\beta_I)) = \frac{1}{2} \text{Tr} \left\{ \sqrt{(\rho_S(t) - \rho_S^{\text{th}}(\beta_I))^2} \right\} \quad (1)$$

- ▶ Exponential dependence with the coupling strength between environments.
- ▶ When the temperatures between reservoirs are closer, it becomes larger.





## Result - Prethermalization time

Trace distance between the prethermal state and the evolved state of the system to study dependence of prethermalization time on other parameters.

$$T(\rho_S(t), \rho_S^{\text{th}}(\beta_I)) = \frac{1}{2} \text{Tr} \left\{ \sqrt{(\rho_S(t) - \rho_S^{\text{th}}(\beta_I))^2} \right\} \quad (1)$$

- ▶ Exponential dependence with the coupling strength between environments.
- ▶ When the temperatures between reservoirs are closer, it becomes larger.
- ▶ Hotter RI yields longer prethermalization times, as well as colder RI.

# Results - Multiple environments



# Concluding Remarks

- ▶ New model to include environments that are not in direct contact with an open quantum system

## Concluding Remarks

- ▶ New model to include environments that are not in direct contact with an open quantum system
- ▶ We extended the usual methods used to study OQS to explore this more complex scenario.

## Concluding Remarks

- ▶ New model to include environments that are not in direct contact with an open quantum system
- ▶ We extended the usual methods used to study OQS to explore this more complex scenario.
- ▶ This model allows to indirectly control the asymptotic state of a system by modifying an environment that is not in direct contact with it

## Concluding Remarks

- ▶ New model to include environments that are not in direct contact with an open quantum system
- ▶ We extended the usual methods used to study OQS to explore this more complex scenario.
- ▶ This model allows to indirectly control the asymptotic state of a system by modifying an environment that is not in direct contact with it
- ▶ Model reminiscent of the layered structure of quantum computers, with different layers that are colder close to the qubits.